

Meso**Description:**

Meso is a viscoelastic gel of non-cross-linked hyaluronic acid, sterile, non pyrogenic, colourless and of nonanimal origin, with almost no chance of side effects. It can effectively protect the epidermis cell, prevent skin aging, moisturize skin and make skin brighter, smooth, and naturally healthy. Each box contains one pre-filled syringes of Meso. For each syringe, the box contains two sterile 30G 1/2" needles and two trackability labels (one to be given to the patient, and the other one to be kept by the doctor in the patient's file). The volume of each syringe is shown on the packaging carton as well as on the syringe.

Composition:

Citrate Buffered Saline
Sodium Hyaluronate (1.5%)

Indication

Meso is injected into dermis to revitalize the skin radiance, moisturize the skin and improve the skin tone.

Mode of action

Hyaluronic acid is a major constituent of ground substance. Its heliocoidal structure enables it to trap one thousand times its weight in water. It is widely distributed in cutaneous tissue and hydrates the skin, though its content diminishes with age. As the skin thins and becomes fragile, the initial skin fracture lines appear. Meso is a hyaluronic acid gel which is injected into the dermis. Thanks to its remarkable fluidity and its visco-elastic properties, Meso diffuses uniformly into cutaneous tissue, maintaining its hydration and reinforcing the dermis. Meso is biodegradable and is absorbed over time. Periodic treatment sessions are necessary to obtain a satisfactory level of skin hydration.

Dosage and Usage

Meso should be injected slowly and superficially over the treatment sites using the multipuncture technique with the aid of the 30G 1/2" needles provided. Manually stretch the treatment sites and carry out multipunctures of 2 to 3 millimetres in diameter separated by one centimetre to a depth of approximately two millimetres. The volume to be injected depends on the size of the area to be treated. It is generally advised to count 2 to 3 treatment sessions at an interval of 3 or 4 weeks, followed by a session every 2 to 3 months in order to maintain the level of hydration.

Precautions and Contraindications

-Normal infection control precautions should be taken as with any other injection.
-Before treatment, a thorough consultation should take place between the dermatologist and the patient, the consultation should cover indications, expected outcomes, potential side effects and complications, anesthesia and aftercare advice.
-Do not administer to patients with hypersensitivity to the product components.
-Do not administer to patients with a history of autoimmune disease or who are receiving immune therapy.

-Do not use in patients presenting an inflammation or irritation of treatment areas.
-Do not inject intravascularly for there is a potential risk of vascular occlusion.
-Do not use in patients with bleeding disorders.

-Do not use in patients who are taking thrombolytics or anticoagulants or have taken inhibitors of platelet aggregation in the preceding 2 weeks.
-Redness, swelling, edema, hematomas, itching, and slight pain at the injection area can occur following the treatment and generally resolve in 72 hours.
-All side effects other than those described above must be reported to the dermatologist, and then inform the product's retailer as soon as possible.
-Special attention should be paid when injected around the eyes.

-People under 18 years are not recommended for use.

-People during a pregnancy or breastfeeding are not recommended for use.

-After use, syringes and needles should be handled as potential biohazards. Disposal should be in accordance with accepted medical practice and applicable local, state and federal requirements.

-Patients should minimize exposure of the treated area to excessive sun, UV lamp exposure and extreme cold weather at least until any initial swelling and redness has resolved.

-Don't use in combination with peeling, laser treatment or ultrasonic-based treatment.

Warnings

-The product must not be injected into blood vessels. Introduction into the vasculature may lead to embolization, occlusion of the vessels, ischemia, or infarction.
-Verify the integrity of the product and expiry date before use.
-Do not use after the expiration date.
-Single use only. Do not resterilize. Do not use if package is damaged or opened.
-The expiration date is valid when product is stored properly in its original package.
-Not for intravenous or intramuscular use.
-Do not mix with other products before injection.
-Keep out of reach of children.
-Product must be injected into non-inflamed, disinfected, healthy skin.

-In order to minimize the risks of potential complications, this product should only be used by health care practitioners who have appropriate training, experience, and who are knowledgeable about the anatomy at and around the site of injection.

-Do not administer to patients with a history of autoimmune disease or who are receiving immune therapy.
-Do not use in patients presenting an inflammation or irritation of treatment areas.
-Do not use in patients with bleeding disorders.
-Do not use in patients with bleeding disorders.

-Do not use in patients who are taking thrombolytics or anticoagulants or have taken inhibitors of platelet aggregation in the preceding 2 weeks.
-Redness, swelling, edema, hematomas, itching, and slight pain at the injection area can occur following the treatment and generally resolve in 72 hours.
-All side effects other than those described above must be reported to the dermatologist and then inform the product's retailer as soon as possible.

-Special attention should be paid when injected around the eyes.

-People under 18 years are not recommended for use.

-People during a pregnancy or breastfeeding are not recommended for use.

-After use, syringes and needles should be handled as potential biohazards. Disposal should be in accordance with accepted medical practice and applicable local, state and federal requirements.

-Patients should minimize exposure of the treated area to excessive sun, UV lamp exposure and extreme cold weather at least until any initial swelling and redness has resolved.

-Don't use in combination with peeling, laser treatment or ultrasonic-based treatment.

Hair Filler**Description:**

Hair Filler is an effective and innovative hair growth treatment made from most advanced formulation to combat thinning hair, it enhances the physical appearance and causes rejuvenation for thicker, fuller and healthier hair. The product formulated with a complex blend of peptides has been proven to enhance anchoring of the hair, increase hair growth and hair follicle size while reducing apoptosis (programmed cell death) that causes follicular miniaturization, help strengthen already existing hair while stimulating growth in areas that are lacking thickness. If your hair is thinning and you wish to have thicker, fuller hair, this is the serum for you.

Composition:

Phosphate Buffered Saline
Sodium Hyaluronate (0.8%)
Copper Peptide (>20 ppm)
Decapeptide-18 (>20 ppm)
Biotinyl Tripeptide-1 (>20 ppm)
Acetyl Tetrapeptide-3 (>20 ppm)

Indication

Hair Filler is injected into dermis at the site of the scalp area. It helps with hair loss prevention and hair regeneration.

Dosage and Usage

-Determine the accurate area of injection, clean the injection area.
-Injection Depth: Superficial to mid dermis
-Injection Point : 0.2-0.3 cm per line
-Injection Dosage: 0.02-0.05 ml per point
-1 session every 2 weeks for 12 weeks
-Regular maintenance sessions (approximately every 1 month) occur as planned with your practitioner ensuring lasting results.

Precautions and Contraindications

-Normal infection control precautions should be taken as with any other injection.
-Before treatment, a thorough consultation should take place between the dermatologist and the patient, the consultation should cover indications, expected outcomes, potential side effects and complications, anesthesia and aftercare advice.
-Do not administer to patients with hypersensitivity to the product components.
-Do not administer to patients with a history of autoimmune disease or who are receiving immune therapy.
-Do not use in patients presenting an inflammation or irritation of treatment areas.
-Do not use in patients with bleeding disorders.
-Do not use in patients with bleeding disorders.

-Do not use in patients who are taking thrombolytics or anticoagulants or have taken inhibitors of platelet aggregation in the preceding 2 weeks.
-Redness, swelling, edema, hematomas, itching, and slight pain at the injection area can occur following the treatment and generally resolve in 72 hours.
-All side effects other than those described above must be reported to the dermatologist and then inform the product's retailer as soon as possible.

-Special attention should be paid when injected around the eyes.

-People under 18 years are not recommended for use.

-People during a pregnancy or breastfeeding are not recommended for use.

-After use, syringes and needles should be handled as potential biohazards. Disposal should be in accordance with accepted medical practice and applicable local, state and federal requirements.

-Patients should minimize exposure of the treated area to excessive sun, UV lamp exposure and extreme cold weather at least until any initial swelling and redness has resolved.

-Don't use in combination with peeling, laser treatment or ultrasonic-based treatment.

retailer as soon as possible.

-People under 18 years are not recommended for use.

-People during a pregnancy or breastfeeding are not recommended for use.

-After use, syringes and needles should be handled as potential biohazards. Disposal should be in accordance with accepted medical practice and applicable local, state and federal requirements.

-Patients should minimize exposure of the treated area to excessive sun, UV lamp exposure and extreme cold weather at least until any initial swelling and redness has resolved.

Warnings

-The product must not be injected into blood vessels. Introduction into the vasculature may lead to embolization, occlusion of the vessels, ischemia or infarction.
-Verify the integrity of the product and expiry date before use.
-Do not use after the expiration date.
-Single use only. Do not resterilize. Do not use if package is damaged or opened.
-The expiration date is valid when product is stored properly in its original package.
-Not for intravenous or intramuscular use.
-Do not mix with other products before injection.
-Keep out of reach of children.
-Product must be injected into non-inflamed, disinfected, healthy skin.

-In order to minimize the risks of potential complications, this product should only be used by health care practitioners who have appropriate training, experience, and who are knowledgeable about the anatomy at and around the site of injection.

Storage Condition:

Store at 2 C -30 C , protect from sunlight. Do not freeze. Do not use after the expiration date.

Shelf Life:

3 years under recommended condition.

Lift**Description:**

Lift is an effective and innovative treatment used in adults to improve the appearance of facial sag. The active ingredient is DMAE, a naturally occurring substance that facilitates the synthesis of a neurotransmitter acetylcholine, a chemical that creates muscle tone. More muscle tone beneath the skin results in tighter, smoother skin. DMAE acts immediately to improve the appearance of sagging and flaccid skin. Its firming action works by protecting the elastin and collagen cross-links. Lift provides an instant face or body lift effect, while defending the skin against free radicals and simultaneously infusing it with moisture. Lift can help slow the progress of wrinkles and sagging and then greatly improve the appearance of wrinkles and sagging in the face.

Composition:

Phosphate Buffered Saline
Sodium Hyaluronate (0.8%)
Arginine (>20 ppm)
DMAE (0.5%)

Indication

Lift is injected into dermis layer for improvement in the appearance of wrinkles and sagging in adults.

Dosage and Usage

-Determine the accurate area of injection, clean the injection area.
-Injection Depth: Superficial to mid dermis
-Injection Point : 0.2-0.3 cm per line
-Injection Dosage: 0.02-0.05 ml per point
-1 session every 2 weeks for 8 weeks
-Regular maintenance sessions (approximately every 2 months) occur as planned with your practitioner ensuring lasting results.

Precautions and Contraindications

-Normal infection control precautions should be taken as with any other injection.
-Before treatment, a thorough consultation should take place between the dermatologist and the patient, the consultation should cover indications, expected outcomes, potential side effects and complications, anesthesia and aftercare advice.
-Do not administer to patients with hypersensitivity to the product components.
-Do not administer to patients with a history of autoimmune disease or who are receiving immune therapy.
-Do not use in patients presenting an inflammation or irritation of treatment areas.

-Do not use in patients with bleeding disorders.
-Do not use in patients with bleeding disorders.
-Do not use in patients who are taking thrombolytics or anticoagulants or have taken inhibitors of platelet aggregation in the preceding 2 weeks.
-Redness, swelling, edema, hematomas, itching, and slight pain at the injection area can occur following the treatment and generally resolve in 72 hours.
-All side effects other than those described above must be reported to the dermatologist, and then inform the product's retailer as soon as possible.

-Special attention should be paid when injected around the eyes.

-People under 18 years are not recommended for use.

-People during a pregnancy or breastfeeding are not recommended for use.

-After use, syringes and needles should be handled as potential biohazards. Disposal should be in accordance with accepted medical practice and applicable local, state and federal requirements.

-Patients should minimize exposure of the treated area to excessive sun, UV lamp exposure and extreme cold weather at least until any initial swelling and redness has resolved.

-Don't use in combination with peeling, laser treatment or ultrasonic-based treatment.

Lift**Description:**

Lift is an effective and innovative treatment used in adults to improve the appearance of facial sag. The active ingredient is DMAE, a naturally occurring substance that facilitates the synthesis of a neurotransmitter acetylcholine, a chemical that creates muscle tone. More muscle tone beneath the skin results in tighter, smoother skin. DMAE acts immediately to improve the appearance of sagging and flaccid skin. Its firming action works by protecting the elastin and collagen cross-links. Lift provides an instant face or body lift effect, while defending the skin against free radicals and simultaneously infusing it with moisture. Lift can help slow the progress of wrinkles and sagging and then greatly improve the appearance of wrinkles and sagging in the face.

Composition:

Phosphate Buffered Saline
Sodium Hyaluronate (0.8%)
Arginine (>20 ppm)
DMAE (0.5%)

Indication

Lift is injected into dermis layer for improvement in the appearance of wrinkles and sagging in adults.

Dosage and Usage

-Determine the accurate area of injection, clean the injection area.
-Injection Depth: Superficial to mid dermis
-Injection Point : 0.2-0.3 cm per line
-Injection Dosage: 0.02-0.05 ml per point
-1 session every 2 weeks for 8 weeks
-Regular maintenance sessions (approximately every 2 months) occur as planned with your practitioner ensuring lasting results.

Precautions and Contraindications

-Normal infection control precautions should be taken as with any other injection.
-Before treatment, a thorough consultation should take place between the dermatologist and the patient, the consultation should cover indications, expected outcomes, potential side effects and complications, anesthesia and aftercare advice.
-Do not administer to patients with hypersensitivity to the product components.
-Do not administer to patients with a history of autoimmune disease or who are receiving immune therapy.
-Do not use in patients presenting an inflammation or irritation of treatment areas.

-Do not use in patients with bleeding disorders.
-Do not use in patients with bleeding disorders.
-Do not use in patients who are taking thrombolytics or anticoagulants or have taken inhibitors of platelet aggregation in the preceding 2 weeks.
-Redness, swelling, edema, hematomas, itching, and slight pain at the injection area can occur following the treatment and generally resolve in 72 hours.
-All side effects other than those described above must be reported to the dermatologist, and then inform the product's retailer as soon as possible.

-Special attention should be paid when injected around the eyes.

-People under 18 years are not recommended for use.

-People during a pregnancy or breastfeeding are not recommended for use.

-After use, syringes and needles should be handled as potential biohazards. Disposal should be in accordance with accepted medical practice and applicable local, state and federal requirements.

-Patients should minimize exposure of the treated area to excessive sun, UV lamp exposure and extreme cold weather at least until any initial swelling and redness has resolved.

-Don't use in combination with peeling, laser treatment or ultrasonic-based treatment.

Lift**Description:**

Lift is an effective and innovative treatment used in adults to improve the appearance of facial sag. The active ingredient is DMAE, a naturally occurring substance that facilitates the synthesis of a neurotransmitter acetylcholine, a chemical that creates muscle tone. More muscle tone beneath the skin results in tighter, smoother skin. DMAE acts immediately to improve the appearance of sagging and flaccid skin. Its firming action works by protecting the elastin and collagen cross-links. Lift provides an instant face or body lift effect, while defending the skin against free radicals and simultaneously infusing it with moisture. Lift can help slow the progress of wrinkles and sagging and then greatly improve the appearance of wrinkles and sagging in the face.

Composition:

Phosphate Buffered Saline
Sodium Hyaluronate (0.8%)
Arginine (>20 ppm)
DMAE (0.5%)

Indication

Lift is injected into dermis layer for improvement in the appearance of wrinkles and sagging in adults.

Dosage and Usage

-Determine the accurate area of injection, clean the injection area.
-Injection Depth: Superficial to mid dermis
-Injection Point : 0.2-0.3 cm per line
-Injection Dosage: 0.02-0.05 ml per point
-1 session every 2 weeks for 8 weeks
-Regular maintenance sessions (approximately every 2 months) occur as planned with your practitioner ensuring lasting results.

Precautions and Contraindications

-Normal infection control precautions should be taken as with any other injection.
-Before treatment, a thorough consultation should take place between the dermatologist and the patient, the consultation should cover indications, expected outcomes, potential side effects and complications, anesthesia and aftercare advice.
-Do not administer to patients with hypersensitivity to the product components.
-Do not administer to patients with a history of autoimmune disease or who are receiving immune therapy.
-Do not use in patients presenting an inflammation or irritation of treatment areas.

-Do not use in patients with bleeding disorders.
-Do not use in patients with bleeding disorders.
-Do not use in patients who are taking thrombolytics or anticoagulants or have taken inhibitors of platelet aggregation in the preceding 2 weeks.
-Redness, swelling, edema, hematomas, itching, and slight pain at the injection area can occur following the treatment and generally resolve in 72 hours.
-All side effects other than those described above must be reported to the dermatologist, and then inform the product's retailer as soon as possible.

-Special attention should be paid when injected around the eyes.

-People under 18 years are not recommended for use.

-People during a pregnancy or breastfeeding are not recommended for use.

-After use, syringes and needles should be handled as potential biohazards. Disposal should be in accordance with accepted medical practice and applicable local, state and federal requirements.

-Patients should minimize exposure of the treated area to excessive sun, UV lamp exposure and extreme cold weather at least until any initial swelling and redness has resolved.

-Don't use in combination with peeling, laser treatment or ultrasonic-based treatment.

Lift**Description:**

Lift is an effective and innovative treatment used in adults to improve the appearance of facial sag. The active ingredient is DMAE, a naturally occurring substance that facilitates the synthesis of a neurotransmitter acetylcholine, a chemical that creates muscle tone. More muscle tone beneath the skin results in tighter, smoother skin. DMAE acts immediately to improve the appearance of sagging and flaccid skin. Its firming action works by protecting the elastin and collagen cross-links. Lift provides an instant face or body lift effect, while defending the skin against free radicals and simultaneously infusing it with moisture. Lift can help slow the progress of wrinkles and sagging and then greatly improve the appearance of wrinkles and sagging in the face.

Composition:

Phosphate Buffered Saline
Sodium Hyaluronate (0.8%)
Arginine (>20 ppm)
DMAE (0.5%)

Indication

Lift is injected into dermis layer for improvement in the appearance of wrinkles and sagging in adults.

Dosage and Usage

-Determine the accurate area of injection, clean the injection area.
-Injection Depth: Superficial to mid dermis
-Injection Point : 0.2-0.3 cm per line
-Injection Dosage: 0.02-0.05 ml per point
-1 session every 2 weeks for 8 weeks
-Regular maintenance sessions (approximately every 2 months) occur as planned with your practitioner ensuring lasting results.

Precautions and Contraindications

-Normal infection control precautions should be taken as with any other injection.
-Before treatment, a thorough consultation should take place between the dermatologist and the patient, the consultation should cover indications, expected outcomes, potential side effects and complications, anesthesia and aftercare advice.
-Do not administer to patients with hypersensitivity to the product components.
-Do not administer to patients with a history of autoimmune disease or who are receiving immune therapy.
-Do not use in patients presenting an inflammation or irritation of treatment areas.

-Do not use in patients with bleeding disorders.
-Do not use in patients with bleeding disorders.
-Do not use in patients who are taking thrombolytics or anticoagulants or have taken inhibitors of platelet aggregation in the preceding 2 weeks.
-Redness, swelling, edema, hematomas, itching, and slight pain at the injection area can occur following the treatment and generally resolve in 72 hours.
-All side effects other than those described above must be reported to the dermatologist, and then inform the product's retailer as soon as possible.

-Special attention should be paid when injected around the eyes.

-People under 18 years are not recommended for use.

-People during a pregnancy or breastfeeding are not recommended for use.

-After use, syringes and needles should be handled as potential biohazards. Disposal should be in accordance with accepted medical practice and applicable local, state and federal requirements.

-Patients should minimize exposure of the treated area to excessive sun, UV lamp exposure and extreme cold weather at least until any initial swelling and redness has resolved.

-Don't use in combination with peeling, laser treatment or ultrasonic-based treatment.

Lift**Description:**

Lift is an effective and innovative treatment used in adults to improve the appearance of facial sag. The active ingredient is DMAE, a naturally occurring substance that facilitates the synthesis of a neurotransmitter acetylcholine, a chemical that creates muscle tone. More muscle tone beneath the skin results in tighter, smoother skin. DMAE acts immediately to improve the appearance of sagging and flaccid skin. Its firming action works by protecting the elastin and collagen cross-links. Lift provides an instant face or body lift effect, while defending the skin against free radicals and simultaneously infusing it with moisture. Lift can help slow the progress of wrinkles and sagging and then greatly improve the appearance of wrinkles and sagging in the face.

Composition:

Phosphate Buffered Saline
Sodium Hyaluronate (0.8%)
Arginine (>20 ppm)
DMAE (0.5%)

Indication

Lift is injected into dermis layer for improvement in the appearance of wrinkles and sagging in adults.

Dosage and Usage

-Determine the accurate area of injection, clean the injection area.
-Injection Depth: Superficial to mid dermis
-Injection Point : 0.2-0.3 cm per line
-Injection Dosage: 0.02-0.05 ml per point
-1 session every 2 weeks for 8 weeks
-Regular maintenance sessions (approximately every 2 months) occur as planned with your practitioner ensuring lasting results.

Precautions and Contraindications

-Normal infection control precautions should be taken as with any other injection.
-Before treatment, a thorough consultation should take place between the dermatologist and the patient, the consultation should cover indications, expected outcomes, potential side effects and complications, anesthesia and aftercare advice.
-Do not administer to patients with hypersensitivity to the product components.
-Do not administer to patients with a history of autoimmune disease or who are receiving immune therapy.
-Do not use in patients presenting an inflammation or irritation of treatment areas.

-Do not use in patients with bleeding disorders.
-Do not use in patients with bleeding disorders.
-Do not use in patients who are taking thrombolytics or anticoagulants or have taken inhibitors of platelet aggregation in the preceding 2 weeks.
-Redness, swelling, edema, hematomas, itching, and slight pain at the injection area can occur following the treatment and generally resolve in 72 hours.
-All side effects other than those described above must be reported to the dermatologist, and then inform the product's retailer as soon as possible.

-Special attention should be paid when injected around the eyes.

-People under 18 years are not recommended for use.

-People during a pregnancy or breastfeeding are not recommended for use.

-After use, syringes and needles should be handled as potential biohazards. Disposal should be in accordance with accepted medical practice and applicable local, state and federal requirements.

-Patients should minimize exposure of the treated area to excessive sun, UV lamp exposure and extreme cold weather at least until any initial swelling and redness has resolved.

-Don't use in combination with peeling, laser treatment or ultrasonic-based treatment.

Lift**Description:**

Lift is an effective and innovative treatment used in adults to improve the appearance of facial sag. The active ingredient is DMAE, a naturally occurring substance that facilitates the synthesis of a neurotransmitter acetylcholine, a chemical that creates muscle tone. More muscle tone beneath the skin results in tighter, smoother skin. DMAE acts immediately to improve the appearance of sagging and flaccid skin. Its firming action works by protecting the elastin and collagen cross-links. Lift provides an instant face or body lift effect, while defending the skin against free radicals and simultaneously infusing it with moisture. Lift can help slow the progress of wrinkles and sagging and then greatly improve the appearance of wrinkles and sagging in the face.

Composition:

Phosphate Buffered Saline
Sodium Hyaluronate (0.8%)
Arginine (>20 ppm)
DMAE (0.5%)

Indication

Lift is injected into dermis layer for improvement in the appearance of wrinkles and sagging in adults.

Dosage and Usage

-Determine the accurate area of injection, clean the injection area.
-Injection Depth: Superficial to mid dermis
-Injection Point : 0.2-0.3 cm per line
-Injection Dosage: 0.02-0.05 ml per point
-1 session every 2 weeks for 8 weeks
-Regular maintenance sessions (approximately every 2 months) occur as planned with your practitioner ensuring lasting results.

Precautions and Contraindications

-Normal infection control precautions should be taken as with any other injection.
-Before treatment, a thorough consultation should take place between the dermatologist and the patient, the consultation should cover indications, expected outcomes, potential side effects and complications, anesthesia and aftercare advice.
-Do not administer to patients with hypersensitivity to the product components.
-Do not administer to patients with a history of autoimmune disease or who are receiving immune therapy.
-Do not use in patients presenting an inflammation or irritation of treatment areas.

-Do not use in patients with bleeding disorders.
-Do not use in patients with bleeding disorders.
-Do not use in patients who are taking thrombolytics or anticoagulants or have taken inhibitors of platelet aggregation in the preceding 2 weeks.
-Redness, swelling, edema, hematomas, itching, and slight pain at the injection area can occur following the treatment and generally resolve in 72 hours.
-All side effects other than those described above must be reported to the dermatologist, and then inform the product's retailer as soon as possible.

-Special attention should be paid when injected around the eyes.

-People under 18 years are not recommended for use.

-People during a pregnancy or breastfeeding are not recommended for use.